

Thermodynamic Bethe ansatz for generalized extensive statistics

Andrei G. Bytsko

*Steklov Mathematics Institute
Fontanka 27, St.Petersburg, 191011, Russia*

Abstract

We investigate properties of the entropy density related to a generalized extensive statistics and derive the thermodynamic Bethe ansatz equation for a system of relativistic particles obeying such a statistics. We investigate the conformal limit of such a system. We also derive a generalized Y-system. The Gentile intermediate statistics and the statistics of γ -ons are considered in detail. In particular, we observe that certain thermodynamic quantities for the Gentile statistics majorize those for the Haldane-Wu statistics. Specifically, for the effective central charges related to affine Toda models we obtain nontrivial inequalities in terms of dilogarithms.

bytsko@pdmi.ras.ru

Research supported in part by INTAS grant 99-01459 and by RFFI grant 99-01-00101

1 Introduction

Although all experimentally observed particles are either bosons or fermions, the general principles of quantum mechanics do not prohibit existence of particles obeying other types of statistics [1]. The motivation to consider an exotic statistics is that it may provide an *effective* description of the dynamics of particles if their interaction is so strong that the available occupancy for a given state depends on the number of particles already present in the state [2]. An exotic statistics can also arise if particles possess a hidden internal degree of freedom that is invisible in the Hamiltonian but that can become dynamically relevant. Finally, an exotic statistics can emerge in description of physical models in one or two spatial dimensions. The reason is that in this case the multi-particle wave function is not necessarily even or odd and particles (anyons) may obey a fractional statistics [3]. Moreover, in low-dimensional theories the exchange statistics of the fields present in a Hamiltonian is not directly related to the exclusion statistics of the corresponding particles. For instance, although particles in the one-dimensional real coupling affine Toda field theories (ATFT) are bosons, one has to impose the Pauli principle on their momenta in the thermodynamic Bethe ansatz analysis in order to obtain correct values of corresponding central charges [4]. Another example is the conformal field theory, where quasi-particle spectra can be constructed with use of an exotic exclusion statistics [5].

The first attempt to introduce an exotic statistics is attributed to G. Gentile [6], who proposed an intermediate statistics, in which at most G indistinguishable particles can occupy a given state. This intermediate statistics interpolates between fermions and bosons, that are recovered for $G = 1$ and $G = \infty$. A decade after Gentile's work H.S. Green introduced the parafermi statistics [7] that also fixes maximal occupancy for a given state. Since then, various aspects of systems governed by the Gentile or parafermi statistics were discussed in the literature, among them possible applications to the elementary particles theory [8] and to the quark confinement problem in the QCD [9].

Many authors discussed thermodynamic properties of an ideal gas obeying the Gentile or parafermi statistics [6, 10, 11]. In the present paper we will study the thermodynamic limit of one-dimensional relativistic integrable field models governed by a Gentile-like statistics. More precisely, the main purpose of this manuscript is to implement systematically a generalized extensive exclusion statistics into the thermodynamic Bethe ansatz (TBA) analysis of systems where the interaction of particles is relativistic, short-range and characterized by a factorizable scattering matrix.

The TBA was originally developed in the seminal papers by Yang and Yang [12] for treating a one-dimensional ideal gas. In later works [13, 14] this technique was adopted to one-dimensional relativistic models. The boundary condition for a many-particle wave function leads to what is commonly referred to as the Bethe ansatz equation, that provides the quantization condition for possible momenta of the system. The TBA constitutes an interface between massive integrable models and conformal field theories. From the TBA one can extract information about the ultraviolet limit of the underlying massive model, in particular, find the corresponding (effective) central charge.

In the derivation of the TBA equation the underlying exclusion statistics is usually taken to be either of bosonic or fermionic type [4, 12, 13]. In the present paper the TBA equation will be derived and studied for significantly more general situation. Namely, let $W(N, n)$ denote the total dimension of the Hilbert space for a system of n indistinguishable particles

that can occupy N different states in the Fock-space. Assume that there exists a function such that its N -th power is a generating function for the dimensions $W(N, n)$, i.e.

$$(f(t))^N = \sum_{k \geq 0} W(N, k) t^k. \quad (1)$$

In the statistical mechanics, if the variable t is understood as fugacity, the sum on the right hand side is the grand partition function of the system. The property of a grand partition function to be an N -th power of a function independent of N is called *extensivity*. As we are going to demonstrate below, this property allows us to develop the TBA analysis for a rather general choice of $f(t)$. Let us stress that for the purpose of deriving the TBA equation even asymptotical extensivity suffices, i.e., equation (1) should hold in the large N limit.

Of course, if the explicit form of $f(t)$ is known, we can use it to simplify the general formulae. For instance, an instructive example is the Gentile statistics of order G which is defined by the following choice of $f(t)$:

$$f_G(t) = 1 + t + t^2 + \dots + t^G. \quad (2)$$

For $G = 1$ and $G = \infty$ this statistics describes ordinary fermions and bosons, respectively. Although $f_G(t)$ is the simplest non-trivial choice of $f(t)$, it has all features of a generalized extensive statistics. Therefore, the Gentile statistics will be our main working example below.

The paper is organized as follows. In Section 2 we describe possible types of a generalized extensive statistics and study properties of the corresponding entropy densities in the thermodynamic limit. The Gentile statistics and the γ -ons statistics are discussed in this context. In Section 3 we compare properties of the Gentile statistics and the Haldane-Wu statistics. In Section 4 we derive the thermodynamic Bethe ansatz equation and the finite-size scaling function for a relativistic multi-particle system in which statistical interaction is governed by a generalized extensive statistics. In Section 5 we derive the Y-system related to such a generalized statistics. In Section 6 we study the ultraviolet limit of the generalized TBA equation and obtain an expression for the corresponding effective central charge. In Section 7 we compute finite-size scaling functions and central charges related to some affine Toda models for the Gentile statistics and the γ -ons statistics and compare them with their counterparts for the Haldane-Wu statistics. Our conclusions are stated in Section 8.

2 Types of extensive statistics and entropy density

Consider a system of n indistinguishable particles that can occupy N different states in the Fock-space. Assume that the particles obey the Gentile statistics of order G , i.e., that each state can be occupied by at most G particles (with G being a positive integer number). There are several combinatorial ways to compute the total dimension of the Hilbert space for such a system. For instance, we first choose m_1 states which are occupied by at least one particle, then among these m_1 states we choose $m_2 \leq m_1$ states which are occupied by at least two particles, etc. This way of counting yields the following expression for the total dimension of the Hilbert space

$$W_G(N, n) = \sum_{N \geq m_1 \geq \dots \geq m_{G-1} \geq 0} C_N^{m_1} C_{m_1}^{m_2} \dots C_{m_{G-2}}^{m_{G-1}} C_{m_{G-1}}^{n-m_1-\dots-m_{G-1}}, \quad (3)$$

where $C_k^m = k!/(m!(k-m)!)$ are binomial coefficients (and $C_k^m = 0$ if $m > k$ or if $m < 0$). Now, let us compute N -th power of the polynomial $f_G(t)$ defined in (2). Thanks to the obvious recursive relation $f_G(t) = 1 + t f_{G-1}(t)$, we can do this by G consecutive applications of the binomial formula. Then it is easy to see that N -th power of $f_G(t)$ is a generating function for the dimensions $W_G(N, n)$, i.e.

$$(f_G(t))^N = \sum_{n=0}^{GN} W_G(N, n) t^n. \quad (4)$$

Thus, the Gentile statistics is a particular case of an extensive statistics.

Let us stress that the order G of the Gentile statistics must be a positive integer. Although we could try to extend it to other values by formal replacement of (2) with

$$f_G(t) = \frac{1 - t^{G+1}}{1 - t}, \quad (5)$$

the resulting function $f_G(t)$ will not satisfy the important positivity property (see below).

Consider now a more general system of n indistinguishable particles that possesses, at least for large N , the extensivity property (1) with

$$f(t) = \sum_{k=0}^d P_k t^k, \quad (6)$$

where d can be a finite positive number or infinity. The Taylor coefficients P_k are fractional dimensions of levels in a single-state Hilbert subspace and can be regarded as probabilities of occupation of the given state by k particles (see related discussions in [15, 16, 17]). The total dimension of the single-state subspace is $f(1)$. It is natural to require *positivity* of the quantities P_k :

$$P_0 = 1 \quad \text{and} \quad P_k \geq 0 \quad \text{for} \quad k \geq 1. \quad (7)$$

The first condition here implies that the vacuum is realized with probability one independently of the size N of the system. Furthermore, we assume that the series (6) converges in the complex plane for $0 \leq |z| < R$, where $R \leq \infty$. This implies that $f(t)$ belongs to one of the following types:

$$\begin{aligned} \text{I.} & \quad d < \infty, & R &= \infty; \\ \text{II.} & \quad d = \infty, & R &= \infty; \\ \text{IIIa.} & \quad d = \infty, & 1 < R < \infty; \\ \text{IIIb.} & \quad d = \infty, & 0 < R \leq 1. \end{aligned} \quad (8)$$

The types I, II, and III consist of finite degree polynomials, analytic functions with infinite Taylor series, and meromorphic functions, respectively. For example, a finite order Gentile statistics belongs to the type I, the Boltzmann statistics ($f(t) = \exp t$) belongs to the type II, and the bosonic statistics ($f(t) = (1-t)^{-1}$) is of the type IIIb. The reason we divided the type III into two subtypes will become clear below.

Let us remark that, in addition to (7), one may require that $P_1 = 1$ with the motivation that the statistics should not be deformed if only one particle is present in the system. Although all the concrete cases which we consider below do fulfill this requirement, it does not seem to be crucial for the general considerations.

The entropy of the system under consideration is $S = k \ln W(N, n)$, where k is the Boltzmann constant. We need to compute the entropy in the thermodynamic limit, i.e.,

when $N \rightarrow \infty$ and n/N is fixed. The extensivity property (1) allows us to express $W(N, n)$ as the following contour integral

$$W(N, n) = \frac{1}{2\pi i} \oint_{|z|=\rho} dz (f(z))^N z^{-n-1} = \frac{1}{2\pi i} \oint_{|z|=\rho} \frac{dz}{z} \exp\{N(\ln f(z) - \mu \ln z)\}, \quad (9)$$

where $\rho < R$, and we denoted $\mu = n/N$. Now we can find an asymptotic expression for the entropy in the thermodynamic limit by application of the saddle point method to the integral (9). The saddle point of the exponent in (9), denote it x , is found as the positive root of the equation (prime stands for a derivative)

$$x f'(t)|_{t=x} = \mu f(x). \quad (10)$$

Choosing $\rho = x$ in (9), so that the saddle point belongs to the integration contour, we apply the standard result (see, e.g., [18]) for the Laplace integral:

$$W(N, \mu N) = \frac{\exp\{N(\ln f(x) - \mu \ln x)\}}{\sqrt{2\pi N h_\mu(x)}} \left(\frac{1}{x} + O(1/N) \right) \quad \text{as } N \rightarrow \infty, \quad (11)$$

where $h_\mu(t) = f''(t)/(f(t)) + (1 - \mu)/t^2$. For any $f(t)$ satisfying (7) and x obeying (10) we have $h_\mu(x) > 0$ due to the Hadamard theorem [18, 19]. From (11) we obtain the entropy density $s(\mu)$ for a given value of μ

$$\lim_{N \rightarrow \infty} \frac{1}{N} \ln W(N, \mu N) \equiv s(\mu) = \ln f(x) - \mu \ln x, \quad (12)$$

where x is positive and satisfies (10).

Two remarks are in order here. Strictly speaking, if $f(t)$ is degenerated, i.e., if there exists integer $p > 1$ such that $f(t) = \sum_k P_{pk} t^{pk}$, then the above derivation modifies since $f(t)$ has p saddle points. However, the corresponding physical interpretation is simply that particles can be added only in p -tuples. Therefore, it is natural to introduce cluster variables $t' = pt$, $n' = pn$, etc. In terms of these new variables formulae (9)-(11) remain valid.

More important remark is that for deriving (12) the property $P_k \geq 0$ is crucial. In general, without the positivity property (9) does not have a real and uniform in μ limit. Moreover, if we allow $f(t)$ to have negative Taylor coefficients, then some $W(N, n)$ in the expansion of $(f(t))^N$ may also be negative. This leads to an immediate problem with the definition of the entropy as a logarithm of $W(N, n)$. However, if the first negative coefficient in the expansion of $(f(t))^N$ appears only at sufficiently large power n_0 (of order $O(N)$), then $f(t)$ violating the positivity condition can still have well defined entropy density for certain range of μ . For instance, such is the case of the Haldane-Wu statistics with statistical interaction g (see Section 3), where $n_0 \approx N/g$. However, in the present paper we will restrict our consideration, for simplicity, only to partition function strictly obeying the positivity condition.

The entropy density is a key quantity for developing the TBA analysis. From the definition of μ it is clear that $0 \leq \mu \leq d$ for $f(t)$ of the type I and $0 \leq \mu \leq \infty$ for the other types. Formally this follows from equation (10) which shows that μ varies from $\mu(x=0)$ to $\mu(x=R)$. Moreover, $\mu(x)$ is a strictly increasing function because $\partial_x \mu = x \mu h_\mu(x) > 0$ for $x > 0$.

Since $f(t)$ is regular at zero, it is easy to see that $\mu \ln x$ tends to zero for small μ . Together with the condition $P_0 = 1$ this yields $s(0) = 0$. In order to analyze further behaviour of $s(\mu)$,

we employ (10) and derive from (12) that

$$\partial_\mu s(\mu) = -\ln x. \quad (13)$$

Since $\partial_x \mu > 0$, we conclude that $\partial_\mu^2 s(\mu) < 0$, that is $s(\mu)$ is a convex up function. Furthermore, (13) implies that $\max s(\mu) = \ln f(\min\{1, R\})$. Note, however, that $s(\mu)$ is not guaranteed to be positive for the whole range of μ . More precisely, $s(\mu)$ may have one root at the interval $\mu(1) < \mu \leq \mu(R)$. Indeed, for $f(t)$ of the type I we derive from (10) and (12) that

$$s(d) = \ln P_d. \quad (14)$$

That is, $s(\mu)$ is non-negative everywhere at $0 \leq \mu \leq d$ only if the last Taylor coefficient P_d is greater or equal to one. For the other types of $f(t)$ we have the following properties

$$s(\infty) = \infty \quad \text{for type IIIb}, \quad (15)$$

$$s(\infty) = -\infty \quad \text{for types II and IIIa}. \quad (16)$$

Indeed, in the first case $x \leq R \leq 1$, hence, according to (13), $s(\mu)$ is strictly increasing. In the second case we have $s(\infty) = \ln f(1) - \int_{\mu(1)}^\infty d\mu \ln x$. The integral here apparently diverges since it exceeds $\int_{\mu(R_0)}^\infty d\mu \ln R_0$, where we can choose any R_0 such that $1 < R_0 < R$.

Investigation of thermodynamic properties of models in which the entropy density becomes negative is rather problematic. Therefore eqs. (14)-(16) suggest that we should restrict the choice of $f(t)$ to the type I with $P_d \geq 1$ (the Gentile-like statistics) and to the type IIIb (the bose-like statistics). However, let us notice that in the TBA framework the variable x depends on the rapidity θ . For some relativistic models with factorizable scattering the typical picture [4] is that $x(\theta)$ falls off as $|\theta|$ grows (for instance, $x(\theta) = \exp\{-mr \cosh \theta\}$ for the trivial S-matrix). In this case $x(\theta) < x(0)$ and we can consider $f(t)$ of the types II and IIIa if $s(\mu)$ remains non-negative for $\mu \leq \mu(x(0))$.

In the context of this discussion it is instructive to consider the so-called γ -ons [20, 11]. These are particles with statistics interpolating between fermions ($\gamma = 1$) and bosons ($\gamma = -1$) in the following simple way

$$\mu(x) = \frac{x}{1 + \gamma x}. \quad (17)$$

Solving equation (10) for this choice of μ it is easy to find the corresponding single-state partition function:

$$f_\gamma(t) = (1 + \gamma t)^{1/\gamma} = 1 + t + \sum_{k \geq 2} [k-1]_\gamma \frac{t^k}{k!}, \quad (18)$$

where $[k]_\gamma = \prod_{m=1}^k (1 - m\gamma)$. For $\gamma > 0$ the positivity condition (7) is fulfilled only if $\gamma = 1/d$, with d positive integer. In this case $f_\gamma(t)$ is of the type I, the corresponding maximal occupancy is $\mu = d$. Since $\ln P_d < 0$ for $d > 1$, the entropy density becomes negative for certain range of μ . The case of $\gamma = 0$ describes the Boltzmann statistics. Here $f_\gamma(t) = \exp t$ belongs to the type II and $s(\mu)$ is negative for $\mu > e$. Finally, for $\gamma < 0$ the positivity condition (7) is always fulfilled and $f_\gamma(t)$ is of the type IIIa or IIIb depending on the value of γ .

Returning to the Gentile statistics, we can summarize the properties of the corresponding entropy density $s_G(\mu)$ as follows (see Fig. 1 for illustration). $s_G(\mu)$ is a convex up function defined for $0 \leq \mu \leq G$ and vanishing at the end points of this interval. It attains the maximum at $\mu = G/2$:

$$s_G(G/2) = \ln(G + 1). \quad (19)$$

Furthermore, $s_G(\mu)$ possesses the following symmetry

$$s_G(G - \mu) = s_G(\mu). \quad (20)$$

This is a consequence of the equality $W_G(N, GN - n) = W_G(N, n)$ which, in turn, follows from the relation (4).

3 Comparison with Haldane-Wu statistics

It is interesting to compare thermodynamic properties of a system of particles obeying the Gentile statistics with those of a system obeying the so-called Haldane-Wu statistics. For the latter, the total dimension of the Hilbert space is postulated to be [21]

$$\hat{W}_g(N, n) = \frac{(N + (1 - g)n + g - 1)!}{n!(N - gn + g - 1)!}. \quad (21)$$

This expression interpolates between the bosonic ($g = 0$) and the fermionic ($g = 1$) statistics. Introduction of such an interpolating statistics was motivated by Haldane's generalization [2] of the Pauli exclusion principle to the form

$$\Delta D / \Delta n = -g. \quad (22)$$

Here D is the number of available states (holes) before the n -th particle has been added to the system. The parameter g is called the statistical interaction. Properties of an ideal gas obeying the Haldane-Wu statistics were actively discussed in the literature [15, 21, 22]. The corresponding TBA equation for relativistic integrable models was obtained in [23] and investigated in [23, 24].

A direct comparison of the quantities $W_G(N, n)$ and $\hat{W}_g(N, n)$ is problematic if the system contains a finite number of particles. Indeed, formula (3) has a bulky form for generic G (i.e., for $2 \leq G < \infty$). Moreover, it requires additional conventions to make exact sense of expression (21) for generic g (i.e., for $0 < g < 1$). Actually, there exists no prescription for counting of states on the microscopical level that would lead to (21) (see related discussions in [15, 16, 17]). To overcome these difficulties we can compare the two statistics in the thermodynamic limit. More precisely, let us compare the entropy density $s_G(\mu)$ with its counterpart $\hat{s}_g(\mu) \equiv \lim_{N \rightarrow \infty} \frac{1}{N} \ln \hat{W}_g(N, \mu N)$ that is readily derived from (21) with the help of the Stirling formula

$$\hat{s}_g(\mu) = (1 + \mu(1 - g)) \ln(1 + \mu(1 - g)) - \mu \ln \mu - (1 - g\mu) \ln(1 - g\mu). \quad (23)$$

Let us notice that the Haldane-Wu statistics is extensive but the corresponding function $f_g(t)$ does not satisfy the positivity property (7). Indeed, we can reconstruct $f_g(t)$ first as a function of μ by the following formula

$$f(\mu) = \exp\{s(\mu) - \mu \partial_\mu s(\mu)\}, \quad (24)$$

that follows from (12) and (13). For $\hat{s}_g(\mu)$ given by (23) this yields

$$f_g(\mu) = \frac{1 + (1 - g)\mu}{1 - g\mu}, \quad (25)$$

which is a well-defined function. But reexpressing it in terms of t with the help of (10) we will always obtain a function that breaks the positivity property (7). For instance, for $g = 1/2$ we find (which coincides with the result of [15]) $f_g(t) = 1 + t^2/2 + t\sqrt{1+t^2/4} = 1 + t + t^2/2 + t^3/8 - t^5/128 + \dots$. However, the first negative term in expansion of the N -th power of this series appears for $n = 2N + 3$. This implies that in the large N limit thermodynamic quantities are well-defined for $\mu \leq 2$.

For our purposes we can regard (23) as a *definition* of the Haldane-Wu statistics. Then we have a finite maximal occupancy N/g for a given number of particles N simply because $\hat{s}_g(\mu)$ is defined for $0 \leq \mu \leq 1/g$. Furthermore, $\hat{s}_g(\mu)$ is a convex up function vanishing at the end points of this interval. Thus, we see that properties of the entropy density $\hat{s}_g(\mu)$ are similar to those of a type I extensive statistics and, specifically, to those of the Gentile statistics. Therefore, it is natural to ask how much the behaviour of $s_G(\mu)$ differs from $\hat{s}_g(\mu)$ if $g = 1/G$ (when the corresponding maximal occupancies coincide).

First, notice that the function $\hat{s}_{1/G}(\mu)$ cannot coincide with $s_G(\mu)$ identically for generic G since $\hat{s}_g(\mu)$ does not possess a symmetry like (20) if $g \neq 0, 1$. Next, we can compare the mid-point value $\hat{s}_{1/G}(G/2) = \frac{G+1}{2} \ln(G+1) - \frac{G}{2} \ln G$ with the corresponding value of $s_G(\mu)$ given by (19). Taking into account the inequalities

$$\begin{aligned} t \ln t &> (t-1) \ln(t+1) & \text{for } t > 1, \\ t \ln t &< (t-1) \ln(t+1) & \text{for } 0 < t < 1, \end{aligned} \quad (26)$$

we establish that $s_G(G/2) > \hat{s}_{1/G}(G/2)$ for $G > 1$. Actually, numerical computations for various values of $G > 1$ show that $s_G(\mu)$ majorizes $\hat{s}_{1/G}(\mu)$ *everywhere* at $0 < \mu < G$,

$$s_G(\mu) > \hat{s}_{1/G}(\mu) \quad \text{for } G > 1. \quad (27)$$

For illustration, the case of $G = 2$ is presented in Fig. 1.

Thus, we see that for generic G the difference between $\hat{s}_{1/G}(\mu)$ and $s_G(\mu)$ is not small if μ is not close to $\mu = 0$ or $\mu = G$. It is instructive to compare these functions also near the end points (in particular, this will provide an additional support to the assertion (27)).

In a vicinity of $\mu = 0$ equation (10) is solved as $x = \mu + (2\delta_{G,1} - 1)\mu^2 + O(\mu^3)$, and we find that $s_G(\mu) = (1 - \ln \mu)\mu + (1/2 - \delta_{G,1})\mu^2 + O(\mu^3)$, where $\delta_{m,n}$ stands for the Kronecker symbol. Comparing this expansion with $\hat{s}_g(\mu) = (1 - \ln \mu)\mu + (1/2 - g)\mu^2 + O(\mu^3)$ which follows from (23), we infer that for small μ functions $\hat{s}_g(\mu)$ and $s_G(\mu)$ take close values (up to the order μ^2) for *any choice* of G and g . Actually, employing (10), it is easy to derive the following expansion for the entropy density of an arbitrary extensive statistics with $f(t)$ satisfying (7):

$$s(\mu) = (P_1 - \ln \mu)\mu + O(\mu^2). \quad (28)$$

Thus, for small μ in the thermodynamic limit, the Haldane-Wu and Gentile statistics are close not only to each other but to any extensive statistics for which two first Taylor coefficients of $f(t)$ are $P_0 = P_1 = 1$. For instance, for small μ the Haldane-Wu and Gentile statistics are close also to the statistics of γ -ons (18).

Consider now $\mu = G - \epsilon$ for small positive ϵ . Then, employing (28) and the symmetry (20), we derive: $s_G(G - \epsilon) = (1 - \ln \epsilon)\epsilon + O(\epsilon^2)$. On the other hand, setting $\mu = 1/g - \epsilon$ in (23), we obtain $\hat{s}_g(1/g - \epsilon) = (1 - 2 \ln g - \ln \epsilon)g\epsilon + O(\epsilon^2)$. We see that in a vicinity of $\mu = G$ the difference between $s_G(\mu)$ and $\hat{s}_{1/G}(\mu)$ is not even of first order in ϵ but of order $\epsilon \ln \epsilon$ (so that the corresponding plots are visibly different here, see Fig. 1).

Summarizing, it appears plausible that the Gentile statistics of order $G > 1$ *majorizes* the Haldane-Wu statistics with parameter $g = 1/G$ and these statistics are *not close* except for small values of $\mu = n/N$.

4 Thermodynamic Bethe ansatz equation

Now we will consider a relativistic multi-particle system of l different species of particles confined to a finite region of the size L . We denote by n_a the number of particles and by N_a the dimension of the Fock-space related to the species a . In the thermodynamic limit both the dimensions of the Fock-space and the size L of the region where the system is confined approach infinity but the ratios n_a/L remain finite [12]. For the fraction of particles with rapidities between $\theta - \Delta\theta/2$ and $\theta + \Delta\theta/2$ it is convenient to introduce densities

$$\Delta N_a = \rho_a(\theta) \Delta\theta L, \quad \Delta n_a = \rho_a^r(\theta) \Delta\theta L. \quad (29)$$

As usual, the rapidity θ parameterizes the two-momentum $P = m(\cosh \theta, \sinh \theta)$ and the total energy of the system is given by

$$E[\rho^r] = L \sum_{a=1}^l \int_{-\infty}^{\infty} d\theta \rho_a^r(\theta) m_a \cosh \theta. \quad (30)$$

According to (29) the variable μ in (12) is now related to the particle densities

$$\mu_a(\theta) = \rho_a^r(\theta) / \rho_a(\theta). \quad (31)$$

Thus, we can express the entropy as the following functional

$$S[\rho, \rho^r] = kL \sum_{a=1}^l \int_{-\infty}^{\infty} d\theta \rho_a(\theta) s(\mu_a) = kL \sum_{a=1}^l \int_{-\infty}^{\infty} d\theta \{ \rho_a(\theta) \ln f_a(x_a(\theta)) - \rho_a^r(\theta) \ln x_a(\theta) \}, \quad (32)$$

where each $x_a(\theta)$ satisfies (10) and hence it is a function of $\rho_a^r(\theta) / \rho_a(\theta)$. The subscript of f_a means that different species may have different single-state partition functions.

In the formulation of the TBA equation for ordinary statistics it is common to introduce the so-called pseudo-energies $\epsilon_a(\theta)$ such that bosonic (upper sign) and fermionic (lower sign) distributions are $\rho_a^r(\theta) / \rho_a(\theta) = (\exp(\epsilon_a(\theta)) \mp 1)^{-1}$. In view of (10), it is natural to define the pseudo-energies for an arbitrary extensive statistics as follows

$$\epsilon_a(\theta) = -\ln x_a(\theta), \quad (33)$$

so that $\rho_a^r(\theta) / \rho_a(\theta) = -\partial_{\epsilon_a(\theta)} \ln f_a(e^{-\epsilon_a(\theta)})$. For instance, the distribution for the Gentile statistics acquires the following form in terms of the pseudo-energies (with a slightly different meaning it was introduced by Gentile [6] for an ideal gas)

$$\frac{\rho_a^r(\theta)}{\rho_a(\theta)} = \frac{\sum_{k=1}^{G_a} k \exp\{(G_a - k)\epsilon_a(\theta)\}}{\sum_{k=0}^{G_a} \exp\{k\epsilon_a(\theta)\}}. \quad (34)$$

According to the fundamental principles of thermodynamics the equilibrium state of a system is found by minimizing the free energy F . Hence, keeping the temperature constant

we obtain the equilibrium condition by minimizing $F[\rho, \rho^r] = E[\rho^r] - TS[\rho, \rho^r]$ with respect to ρ_a^r . The equilibrium condition reads

$$\frac{\delta F}{\delta \rho_a^r} = \frac{\delta E}{\delta \rho_a^r} - T \frac{\delta S}{\delta \rho_a^r} - T \sum_{b=1}^l \frac{\delta S}{\delta \rho_b} \frac{\delta \rho_b}{\delta \rho_a^r} = 0. \quad (35)$$

The admissible momenta in the system are restricted by the boundary conditions. To derive the corresponding quantization equations one takes a particle in the multi-particle wave function on a trip through the whole system [12, 13]. The particle will scatter with all other particles, which yields the following set of equations determining admissible rapidities

$$\exp(iLm_i \sinh \theta_i) \prod_{j \neq i}^N S_{ij}(\theta_i - \theta_j) = \kappa_i, \quad i = 1, \dots, N, \quad (36)$$

where $N = \sum_a N_a$, and $S_{ij}(\theta)$ is a two-particle scattering matrix. The constant phases κ_i on the r.h.s. of (36) may depend on the particle's species; for our purposes their exact values are irrelevant. Taking the logarithmic derivative of (36) and employing densities as in (29), we obtain

$$m_a \cosh \theta + 2\pi \sum_{b=1}^l (\varphi_{ab} * \rho_b^r)(\theta) = 2\pi \rho_a(\theta). \quad (37)$$

Here $(u * v)(\theta) \equiv 1/(2\pi) \int d\theta' u(\theta - \theta') v(\theta')$ stands for the convolution and we introduced as usual the notation $\varphi_{ab}(\theta) = -i\partial_\theta(\ln S_{ab}(\theta))$.

Employing equations (10), (12) and (31), we compute variations of the entropy (32):

$$\delta S / \delta \rho_a^r = -\ln x_a(\theta), \quad \delta S / \delta \rho_a = \ln f_a(x_a(\theta)). \quad (38)$$

Substitution of these relations into the equilibrium condition (35) together with (37) yields the desired thermodynamic Bethe ansatz equations for a system in which statistical interaction between particles of a -th species is described by a single-state partition function f_a ,

$$\frac{1}{kT} m_a \cosh \theta = -\ln x_a(\theta) + \sum_{b=1}^l (\varphi_{ab} * \ln f_b(x_b))(\theta). \quad (39)$$

In general, this set of integral equations cannot be solved analytically. However, one can try to solve these TBA equations numerically with the help of the iteration method (as we will do for some examples in Section 7). For the ordinary statistics and the Haldane-Wu case it is known that such an iterative procedure converges provided that $\varphi_{ab}(\theta)$ falls off sufficiently fast. The most complete proof of this assertion is presented in [24, 25] and is based on the fixed point theorem. Presumably this proof extends to the case of (39) (although some further restrictions on $f_a(t)$ may be required), but we will not discuss this here. Let us notice only that if (39) can be solved iteratively and $\varphi_{ab}(\theta)$ is symmetric in θ (which is always the case for a unitary S-matrix), then $x_a(\theta)$ is also symmetric in θ .

We now substitute the expressions for the total energy (30) and the entropy (32) together with equations (37) and (39) back into the expression for the free energy, and obtain

$$F(T) = -\frac{LkT}{2\pi} \sum_{a=1}^l m_a \int_{-\infty}^{\infty} d\theta \cosh \theta \ln f_a(x_a(\theta)). \quad (40)$$

The relation between the free energy and the finite-size scaling function is well-known to be $c(T) = -6F(T)/(\pi LT^2)$ [26]. As usual we introduce a variable $r = 1/T$ and set now the Boltzmann constant to be one. Then the finite-size scaling function acquires the form

$$c(r) = \sum_{a=1}^l \frac{6m_a r}{\pi^2} \int_0^\infty d\theta \cosh \theta \ln f_a(x_a(\theta)). \quad (41)$$

Here we assumed that $x_a(\theta) = x_a(-\theta)$. Once more for $f(t) = 1 + t$ and $f(t) = 1/(1 - t)$ we recover the well-known expressions for the fermionic and bosonic scaling functions.

It is instructive to compare the above TBA equation (39) and the finite-size scaling function (41) with those for the Haldane-Wu statistics [23]:

$$\frac{1}{kT} m_a \cosh \theta = \ln(1 + y_a(\theta)) + \sum_{b=1}^l (\Phi_{ab} * \ln(1 + y_b^{-1}))(\theta), \quad (42)$$

$$c_g(r) = \sum_{a=1}^l \frac{6m_a r}{\pi^2} \int_0^\infty d\theta \cosh \theta \ln(1 + y_a^{-1}(\theta)). \quad (43)$$

Here $\Phi_{ab}(\theta) = \varphi_{ab}(\theta) - 2\pi g_{ab} \delta(\theta)$, and g is a matrix which appears on the r.h.s. of (22) if we consider a system of several species. As was discussed in [23], the TBA equation (42) leads to an equivalence principle: two systems having the same mass spectra and identical quantities $\Phi_{ab}(\theta)$ are thermodynamically equivalent (as seen from (43), their finite-size scaling functions coincide). Our TBA equation (39) does not possess such a feature. More precisely, for (39) there exists no way to compensate a difference in statistics by a change of $\varphi_{ab}(\theta)$ independent of the explicit form of the S-matrix.

5 Y-systems

For some class of models it is possible to carry out certain manipulations on the TBA equations such that the original integral TBA equations acquire the form of a set of functional equations in new variables Y_a [14]. These functional equations (commonly referred to as Y-systems) have the further virtue that unlike the original TBA equations they *do not* involve the mass spectrum. In the Y -variables certain periodicities in the rapidities are exhibited more clearly. These periodicities may then be utilized in order to express the quantity $Y_a(\theta)$ as a Fourier series, which in turn is useful to find solution of the TBA equations and expand the scaling function as a power series in the scaling parameter r . We will now demonstrate that similar equations may be derived for a multi-particle system in which the dynamical scattering is governed by the scattering matrix related to the ADE-affine Toda field theories and the statistical interaction is of a general type considered above.

Consider the minimal part (i.e., independent of the coupling constant) of the scattering matrix of the ADE-affine Toda field theories. As was shown in [27], upon appropriate choice of CDD-ambiguities, these S-matrices satisfy the identity

$$\varphi_{ab}\left(\theta + \frac{i\pi}{h}\right) + \varphi_{ab}\left(\theta - \frac{i\pi}{h}\right) = \sum_{c=1}^l I_{ac} \varphi_{cb}(\theta) - 2\pi I_{ab} \delta(\theta), \quad (44)$$

where h denotes the Coxeter number, l the rank and I the incidence matrix of the corresponding Lie algebra. It is then straightforward to derive the “Y-system”

$$Y_a\left(\theta + \frac{i\pi}{h}\right) Y_a\left(\theta - \frac{i\pi}{h}\right) = \prod_{b=1}^l \left(z_b(Y_b(\theta))\right)^{I_{ab}}, \quad (45)$$

where $Y_a(\theta) = x_a^{-1}(\theta)$ and

$$z_a(t) = t f_a(t^{-1}). \quad (46)$$

Equations (45) follow upon first adding (39) at $\theta + \frac{i\pi}{h}$ and $\theta - \frac{i\pi}{h}$ and subtracting I times (39) at θ from the sum. Thereafter we employ the fact [28] that the masses of an ADE affine Toda field theory are proportional to the Perron-Frobenius vector of the corresponding Cartan matrix, i.e., $\sum_b C_{ab} m_b = 4 \sin^2(\pi/(2h)) m_a$. Then, with the help of (44), equations (45) follow.

In comparison with (39) equations (45) have already the virtue that they are simple functional equations and do not involve the mass spectrum. For the Gentile statistics we have $z(t) = t + 1 + t^{-1} + \dots + t^{1-G}$. In particular, for $G = 1$ we have $z(Y) = 1 + Y$ and recover the known fermionic Y-system. The bosonic case ($G = \infty$) corresponds to $z(Y) = Y^2/(Y - 1)$. And the case of $G = 2$ seems to be particularly interesting since here $z(Y) = 1 + Y + Y^{-1}$ possesses an additional symmetry $z(Y) = z(Y^{-1})$ or, equivalently, $z(\epsilon_a(\theta)) = z(-\epsilon_a(\theta))$.

6 Ultraviolet limit

In the small r limit the scaling function $c(r)$ becomes the effective central charge of a conformal field theory [26] which describes the ultraviolet limit of a given massive model. That is, $\lim_{r \rightarrow 0} c(r) = c_{\text{eff}} \equiv c - 24h'$, where c is the conformal anomaly and h' is the lowest conformal weight in the corresponding conformal field theory.

In order to evaluate $c(r)$ in the $r \rightarrow 0$ limit we substitute the derivative of (39) into (41), replacing in both equations $\cosh \theta$ by $\frac{1}{2}e^\theta$. Then we integrate the resulting equation by parts, assuming that $\varphi_{ab}(\theta)$ is symmetric and falls off sufficiently fast (typically, $\varphi_{ab}(\theta) = O(e^{-\theta})$ as $\theta \rightarrow \infty$). Next we substitute back equation (39). Finally we change variables from θ to $x(\theta)$, taking into account that $x(\infty) = 0$ (as follows from (39)), and again integrate by parts. All this yields

$$c_{\text{eff}} = \frac{6}{\pi^2} \sum_{a=1}^l c_a, \quad (47)$$

$$c_a = -\frac{1}{2} \ln x_a \ln f_a(x_a) + \int_0^{x_a} \frac{dt}{t} \ln f_a(t). \quad (48)$$

Here we denoted $x_a \equiv x_a(\theta = 0)$. These quantities can be determined from the *constant* TBA equations that follow from (39) when $r \rightarrow 0$ (this derivation requires that $x_a(\theta)$ become constant in a region of θ of order $-\ln r$ when r is small; see more detailed explanations in [4, 25])

$$\ln x_a = - \sum_{b=1}^l N_{ab} \ln f_b(x_b), \quad a = 1, \dots, l, \quad (49)$$

where $2\pi N_{ab} = - \int_{-\infty}^{\infty} d\theta \varphi_{ab}(\theta)$.

As a simple example we consider the statistics of γ -ons (18), assuming for simplicity that γ is the same for all species (generalization to different γ_a is obvious). Here the integral in

(48) can be computed explicitly in terms of the Euler dilogarithm which is defined for $t \leq 1$ as $Li_2(t) = \sum_{k \geq 1} t^k / k^2$. We have

$$\int_0^x \frac{dt}{t} \ln f_\gamma(t) = \frac{1}{\gamma} \int_0^x \frac{dt}{t} \ln(1 + \gamma t) = -\frac{1}{\gamma} \sum_{k=1}^{\infty} \frac{(-\gamma x)^k}{k^2} = -\frac{1}{\gamma} Li_2(-\gamma x)$$

and hence

$$c_a = -\frac{1}{\gamma} Li_2(-\gamma x_a) - \frac{1}{2\gamma} \ln x_a \ln(1 + \gamma x_a), \quad (50)$$

where x_a satisfies (49). Since $Li_2(-t) = \frac{1}{2} Li_2(t^2) - Li_2(t)$, we can rewrite (50) for positive γ employing the Rogers dilogarithm which is defined as $L(t) = Li_2(t) + \frac{1}{2} \ln t \ln(1 - t)$ for $0 \leq t \leq 1$. Then, using the Abel identity $L(t^2) = 2L(t) - 2L(\frac{t}{1+t})$, we obtain

$$c_a = \frac{1}{\gamma} L\left(\frac{\gamma x_a}{1 + \gamma x_a}\right) + \frac{1}{2\gamma} \ln \gamma \ln(1 + \gamma x_a). \quad (51)$$

Notice that this expression does not diverge for small γ ; $\lim_{\gamma \rightarrow 0} c_a = x_a(1 - \frac{1}{2} \ln x_a)$ is the value corresponding to the Boltzmann statistics.

Consider now the Gentile statistics, assuming that the order G is the same for all species. In this case we also can compute the integral in (48) explicitly:

$$\int_0^x \frac{dt}{t} \ln f_G(t) = \int_0^x \frac{dt}{t} \ln\left(\frac{1 - t^{G+1}}{1 - t}\right) = \sum_{k=1}^{\infty} \frac{1}{k^2} \left(x^k - \frac{x^{k(G+1)}}{G+1}\right) = Li_2(x) - \frac{Li_2(x^{G+1})}{G+1},$$

and we see that for the Gentile statistics expression (48) acquires the following nice form involving only the Rogers dilogarithms:

$$c_a = L(x_a) - \frac{1}{G+1} L(x_a^{G+1}). \quad (52)$$

Introducing, in agreement with (33), the constants $\epsilon_a = -\ln x_a$, we obtain from (52) for $G = \infty$ and $G = 1$

$$c_a = L(e^{-\epsilon_a}) \quad \text{and} \quad c_a = L(e^{-\epsilon_a}) - \frac{1}{2} L(e^{-2\epsilon_a}). \quad (53)$$

These are the well-known values for the bosonic and fermionic statistics [4]. Notice that in the latter case one usually derives $c_a = L(\frac{1}{1+e^{\epsilon_a}})$. This is equivalent to our formula (53) due to the Abel identity. In fact, the Abel identity can be used in a similar way for any G of the form $G = 2^m - 1$. In this case we can rewrite (52) as follows

$$c_a = \sum_{k=1}^m \frac{1}{2^{k-1}} L\left(\frac{(x_a)^{2^{k-1}}}{1 + (x_a)^{2^{k-1}}}\right). \quad (54)$$

In this form c_a is a manifestly positive and monotonic function of x_a (the dilogarithm $L(t)$ grows monotonically for $0 \leq t \leq 1$). Actually, this property holds not only for the Gentile statistics but in general case as well. To prove this statement we take a derivative of (48) and use (10) and the formula (12) for the entropy density. This yields

$$\partial_{x_a} c_a = \frac{s(\mu_a(x_a))}{2x_a}. \quad (55)$$

As we discussed in Section 2, we should consider only such models where $s(\mu(x)) > 0$. In this case the r.h.s. of (55) is positive. Hence, c_a is always a monotonically increasing function of x_a (and therefore a monotonically decreasing function of ϵ_a). Moreover, taking into account (7), we infer from (48) that $c_a(0) = 0$, thus we also proved positivity of $c_a(x_a)$. Since these properties are crucial for a physical interpretation of c_a , let us underline that they hold only if the first condition in (7) is fulfilled. Let us notice also that the properties of c_a together with (49) imply that for any model such that $N_{ab} \geq 0$ (which is the case, for instance, for all the ADE-affine Toda models [4]) the value of c_a does not exceed that of the corresponding free model (where all $N_{ab} = 0$ and consequently all $x_a = 1$). Hence, (48) evaluated for $x_a = 1$ yields the upper bound on c_a ,

$$c_a \leq \int_0^1 \frac{dt}{t} \ln f_a(t). \quad (56)$$

Notice, however, that for a statistics of the type IIIb with $R < 1$ we have $x_a < 1$, so that in this case the upper bound on c_a will be lower than (56). From a physical point of view this implies that such a statistics cannot emerge in models with weak interaction and, in particular, in a free model.

It was argued in [4] that the quantity $\frac{6}{\pi^2}c_a$ (for the ordinary statistics) can be interpreted as a massless degree of freedom associated to the a -th species in the massive model. The properties of c_a which we proved above show that the interpretation of $\frac{6}{\pi^2}c_a$ can be extended to the case of a generalized extensive statistics as well. In this context, it is important to remark that for the ordinary statistics (actually, for the Gentile statistics of any order) we have $\frac{6}{\pi^2}c_a \leq 1$ as follows from (52) (recall that $L(1) = Li_2(1) = \pi^2/6$) if all N_{ab} are non-negative. However, for a general statistics the upper bound (56) on $\frac{6}{\pi^2}c_a$ depends on the choice of f_a and may not equal to one. For instance, taking $f_a(t) = (1-t)^{-m}$ with $m > 1$, we obtain $\frac{6}{\pi^2}c_a = m$ for the corresponding free theory. In such a case, the massless degree of freedom of the corresponding free particle exceeds that of a free boson. This can possibly imply that we have to restrict the choice of statistics to such f_a that $c_a \leq \frac{\pi^2}{6}$.

7 Examples related to affine Toda models

7.1 Ising and Klein-Gordon models

The most simple examples which illustrate the features outlined above more concretely are the Ising model (A_1 -minimal affine Toda field theory) with $S(\theta) = -1$ and the Klein-Gordon model with $S(\theta) = 1$. In both cases equation (39) is solved trivially, $x(\theta) = \exp\{-rm \cosh \theta\}$. Then for the Gentile statistics (2) we can compute the entire scaling function (41):

$$\begin{aligned} c_G(r) &= \frac{6rm}{\pi^2} \int_0^\infty d\theta \cosh \theta \left(\ln(1 - e^{-(G+1)rm \cosh \theta}) - \ln(1 - e^{-rm \cosh \theta}) \right) \\ &= \frac{6rm}{\pi^2} \sum_{k=1}^\infty \frac{1}{k} \left(K_1(krm) - K_1((G+1)krm) \right), \end{aligned} \quad (57)$$

where $K_1(t)$ is the modified Bessel function. We depict $c_G(r)$ in Fig. 2, referring to them by a slight abuse of notation as A_1 . One observes that the difference in statistics affects most severely the ultraviolet region. The scaling functions for different G converge relatively fast towards each other in the infrared regime. These appear to be common features of a general deformed statistics.

As seen from Fig. 2, the behaviour of $c_G(r)$ in the ultraviolet region (there is no first order term in the small r expansion of $c_G(r)$) is similar for any finite G to that of the fermionic scaling function and is different from the bosonic case (where the first order term is present). To explain this we observe that, as follows from (57),

$$c_G(r) = c_b(r) - \frac{1}{G+1} c_b((G+1)r), \quad (58)$$

where $c_b(r)$ is the bosonic scaling function. Now it is obvious that the first order terms on the r.h.s. of (58) cancel each other for any finite G .

The well-known property of the asymptotic behaviour of the modified Bessel function, $\lim_{t \rightarrow 0} tK_1(t) = 1$, allows us to derive from (57)

$$c_G^{\text{eff}} = \lim_{r \rightarrow 0} c_G(r) = \frac{6}{\pi^2} \sum_{k=1}^{\infty} \frac{1}{k^2} \left(1 - \frac{1}{G+1}\right) = \frac{G}{G+1}, \quad (59)$$

which is in agreement with (52) since $x_a = 1$ now. It is worth noticing that c_G^{eff} can be identified for every positive integer G as the effective central charge of a minimal conformal model $\mathcal{M}(s, t)$ (such that $st = 6(G+1)$). In particular, $G = 1, 2, 3, 4$ correspond to the $\mathcal{M}(3, 4)$, $\mathcal{M}(2, 9)$, $\mathcal{M}(3, 8)$, $\mathcal{M}(5, 6)$ minimal models, respectively.

For the γ -ons (18) the scaling function related to the A_1 -models can also be computed:

$$c_\gamma(r) = \frac{6rm}{\gamma\pi^2} \int_0^\infty d\theta \cosh \theta \ln(1 + \gamma e^{-rm \cosh \theta}) = \frac{6rm}{\pi^2} \sum_{k=1}^{\infty} \frac{1}{k} (-\gamma)^{k-1} K_1(krm). \quad (60)$$

Using again the asymptotic behaviour of the modified Bessel function, we obtain $c_\gamma^{\text{eff}} = \lim_{r \rightarrow 0} c_\gamma(r) = -\frac{6}{\gamma\pi^2} Li_2(-\gamma)$ in agreement with (50). Here, unlike in the Gentile case, it is not easy to see whether the effective central charge c_γ^{eff} has rational value for a given γ . As we discussed in Section 2, positive γ is to be of the form $\gamma = 1/d$ with d positive integer. It appears then that the only d leading to rational value of c_γ^{eff} is $d = 1$. On the other hand, γ varies continuously in the range $-1 \leq \gamma \leq 0$, so that any rational value of c_γ^{eff} not exceeding $\frac{6}{\pi^2}$ occurs here. For the Boltzmann statistics $c_0^{\text{eff}} = \frac{6}{\pi^2}$ is irrational. Finally, $\gamma < -1$ is the case of a type IIIb statistics with $R < 1$, which, as we discussed above, cannot emerge in a free model.

7.2 Scaling Potts and Yang-Lee models

Next we consider the scaling Potts model, which was studied previously in the TBA framework for the fermionic statistics in [13] and for the Haldane-Wu statistics in [23]. The two particles in the model are conjugate to each other and consequently their masses are the same, $m_1 = m_2 = m$. The conjugate particle occurs as a bound state when two particles of the same species scatter. The S-matrix of the scaling Potts model equals the minimal S-matrix of the A_2 -affine Toda field theory. The corresponding quantities $\varphi_{ab}(\theta)$ that enter the TBA equation are given by

$$\varphi_{11}(\theta) = \varphi_{22}(\theta) = \frac{-\sqrt{3}}{2 \cosh \theta + 1} \quad \text{and} \quad \varphi_{12}(\theta) = \frac{\sqrt{3}}{1 - 2 \cosh \theta}. \quad (61)$$

Consider the scaling Potts model with both species of particles obeying the Gentile statistics of the same order G . In this case the Z_2 -symmetry of the model is preserved so

that $x_1(\theta) = x_2(\theta) \equiv x(\theta)$ and (39) reduces to a single integral equation. It appears to be impossible to find an analytic solution $x(\theta)$ to this equation, but it is straightforward to solve it numerically. Taking $x^{[0]}(\theta) = \exp\{-rm \cosh \theta\}$ as the first approximation, we can iterate (39) as follows

$$\ln(x^{[n+1]}(\theta)) = -rm \cosh \theta - \frac{2\sqrt{3}}{\pi} \int_{-\infty}^{\infty} d\theta' \frac{\cosh(\theta - \theta')}{1 + 2 \cosh 2(\theta - \theta')} \ln(f_G(x^{[n]}(\theta'))). \quad (62)$$

Convergence is achieved relatively quickly (it depends on the value of mr). The results for $G=1$ (which is in complete agreement with the calculation in [13]) and for $G=2$ are shown in Fig. 3. To make contact with the literature, we introduced the quantity $L(\theta) = \ln f_G(x(\theta))$. One observes the typical plateau $L(\theta) = \text{const}$ for some region of θ when $mr \rightarrow 0$, which is required to derive (49). We see also that for the same value of mr the functions $L(\theta)$ corresponding to different G have similar profiles but different heights of the plateau.

Having solved the generalized TBA equation, we can substitute $x(\theta)$ into (41) and compute the entire scaling function $c_G(mr)$. The result of the numerical computation is shown in Fig. 4. The behaviour of the scaling functions is similar to what we have already observed in the A_1 case.

Due to the fact that the scattering matrices of the scaling Potts model and the scaling Yang-Lee model are related as $S^{YL}(\theta) = S_{11}^{A_2}(\theta) S_{12}^{A_2}(\theta)$, the generalized TBA equations for the two models coincide. The only difference is that the scaling Yang-Lee model has only one particle. Therefore its scaling function equals a half of that for the scaling Potts model.

7.3 Comparison with Haldane-Wu statistics

In Section 3 we have seen that the entropy density corresponding to the Gentile statistics majorizes that for the Haldane-Wu statistics. It appears (at least for certain models) that an analogous relation holds for the corresponding finite-size scaling functions. Specifically, if $c_G(r)$ is the scaling function (41) of A_1 or A_2 minimal affine Toda model where all species obey order G Gentile statistics and $c_g(r)$ is the scaling function (43) of the same model where the Haldane-Wu statistical interaction is of the form $g_{ab} = \frac{1}{G} \delta_{ab}$, then for all $r \geq 0$ we have

$$c_G(r) > c_g(r), \quad \text{where} \quad g_{ab} = \frac{1}{G} \delta_{ab} \quad \text{and} \quad G > 1. \quad (63)$$

For the A_1 case we can rigorously prove relation (63) since the solutions of the generalized TBA equations (39) and (42) are found explicitly:

$$g \ln y(\theta) + (1 - g) \ln(1 + y(\theta)) = mr \cosh \theta = -\ln x(\theta). \quad (64)$$

This equation allows us to express $x(\theta)$ in terms of $y(\theta)$:

$$x(\theta) = y^{-g}(\theta) (1 + y(\theta))^{g-1}. \quad (65)$$

Moreover, since $mr \cosh \theta > 0$, we infer from (64) that

$$y(\theta) > y_0, \quad \text{where} \quad y_0^g = (1 + y_0)^{g-1}. \quad (66)$$

As seen from (41) and (43), in order to establish (63) it suffices to show that $f_{G=1/g}(x(\theta)) > 1 + y^{-1}(\theta)$. The latter relation is equivalent to the inequality

$$1 - (1 + y(\theta))^{-\frac{1}{g}} > y^g(\theta) (1 + y(\theta))^{-g}, \quad (67)$$

as can be verified by simple algebraic manipulations using formula (65). Now it is elementary to check that (67) is valid provided that (66) holds. Thus, we proved the assertion (63) in the A_1 case. For illustration, the scaling function $c_g(r)$ for $g=1/2$ Haldane-Wu statistics is shown in Fig. 2. It lies everywhere below $c_G(r)$ for $G=2$ Gentile statistics with the maximal difference reached in the ultraviolet region.

For a higher rank case, already for A_2 , the generalized TBA equations are integral and it is not clear whether we can compare $x_a(\theta)$ and $y_a(\theta)$ analytically. However, numerical computations for various G show that (63) holds in the A_2 case as well. For illustration, we present in Fig. 4 the scaling function $c_g(r)$ for the Haldane-Wu statistics with $g_{ab} = \frac{1}{2}\delta_{ab}$.

A particular consequence of (63) is the following inequality for the corresponding central charges:

$$c_G^{\text{eff}} > c_g^{\text{eff}}, \quad \text{for } g_{ab} = \frac{1}{G}\delta_{ab} \quad \text{and} \quad G > 1. \quad (68)$$

Here c_G^{eff} is found from (47), (49) and (52) whereas c_g^{eff} is the effective central charge related to the Haldane-Wu statistics with $g_{ab} = \frac{1}{G}\delta_{ab}$ and is given by [23]

$$c_g^{\text{eff}} = \frac{6}{\pi^2} \sum_{a=1}^l L\left(\frac{1}{1+y_a}\right), \quad \text{where} \quad \ln(1+y_a) = \sum_{b=1}^l (N_{ab} + g\delta_{ab}) \ln(1+y_a^{-1}). \quad (69)$$

Again, in the A_1 case we can prove relation (68) directly. Here c_G^{eff} has a simple form (59) and therefore (68) reduces to

$$\frac{1}{1+g} > \frac{6}{\pi^2} L\left(\frac{1}{1+y_0}\right), \quad (70)$$

where y_0 is defined as in (66). Notice that $g < y_0 < 1$ for $g \neq 0, 1$ (since $y_0 \leq g$ would contradict (26)). Therefore, $\frac{1}{2} < \frac{1}{1+y_0} < \frac{1}{1+g}$. Using now a property of the dilogarithm,

$$L(t) < \frac{\pi^2}{6}t \quad \text{for } \frac{1}{2} < t < 1; \quad L(t) > \frac{\pi^2}{6}t \quad \text{for } 0 < t < \frac{1}{2}, \quad (71)$$

we establish the desired inequality (70).

As we discussed above, the order G of the Gentile statistics is to be a positive integer. However, if we regard (47), (49) and (52) as a *definition* of the quantity c_G^{eff} , then (70) extends to any $G > 1$ (since the proof does not require that G be an integer) as shown in Fig. 5. Furthermore, considering G in the range $0 < G < 1$, we can prove with the help of (26) that here $y_0 < g$ and hence $\frac{6}{\pi^2}L(\frac{1}{1+y_0}) > \frac{1}{1+g}$ due to (71). Thus, for this range of G we have a *reverse* inequality, i.e.,

$$c_G^{\text{eff}} < c_g^{\text{eff}}, \quad \text{for } g_{ab} = \frac{1}{G}\delta_{ab} \quad \text{and} \quad 0 < G < 1. \quad (72)$$

As we will see below, this inequality holds for higher rank cases as well. But let us stress again that c_G^{eff} in (72) is a formally defined quantity which is not related directly to the Gentile statistics.

The inequality for the central charges (68) is a weaker statement than the inequality for the scaling functions (63). Furthermore, (68) involves dilogarithms, which makes its proof more complicated (already in the A_1 case we used a rather subtle property (71)). But the advantage of (68) for a numerical verification is that it relates numbers and not functions as (63). Moreover, since $c(r)$ is a continuous function, validity of (68) implies that (63) holds at least in some ultraviolet region. Therefore, we will discuss below validity of (68) for

several affine Toda models. As a mathematical by-product, this provides us with interesting dilogarithm inequalities.

In the A_2 case, eq. (68) contains dilogarithms in a more involved way. As seen from (49), (52) and (69), it is equivalent to the following inequality

$$L(\tilde{x}_0) - \frac{g}{1+g} L\left(\tilde{x}_0^{1+\frac{1}{g}}\right) > L\left(\frac{1}{1+\tilde{y}_0}\right), \quad \text{for } 0 < g < 1. \quad (73)$$

Here $\tilde{x}_0$ and $\tilde{y}_0$ are determined from the following equations that follow from (49) and (69) upon noticing that $N_{11} + N_{12} = 1$

$$\tilde{x}_0 f_{1/g}(\tilde{x}_0) = 1 \quad \text{and} \quad \tilde{y}_0^{1+g} = (1 + \tilde{y}_0)^g. \quad (74)$$

It is worth remarking that, as was noticed in [23], $\frac{6}{\pi^2} L\left(\frac{1}{1+\tilde{y}_0}\right)$ coincides with the effective central charge of the Calogero-Sutherland model with the coupling constant $\lambda = g$.

Numerical computations show that (73) indeed holds for any $0 < g < 1$, that is for any $G > 1$ (here again we need not to restrict G to be an integer). Furthermore, like in the A_1 case, inequality (73) reverses for $0 < G < 1$, i.e., for $g > 1$. For illustration, we depict the corresponding c_G^{eff} and c_g^{eff} in Fig. 5. This result provides an additional support to our claim that in the A_2 case the entire scaling function $c_G(r)$ majorizes $c_g(r)$ for $G > 1$.

It is interesting to remark that, as follows from (52) and (69), the bosonic version ($G = \infty$ or, equivalently, $g = 0$) of the A_2 -minimal affine Toda model has $c_a = \frac{1}{2}$. That is, then each species in the scaling Potts model (or the only species in the scaling Yang-Lee model) appears to be a free fermion, whereas the entire central charge is that of a free boson (see also related comments in [23]). That is why all curves on Fig. 5 converge to the same value $c = 1$.

7.4 Higher rank cases

Now we discuss briefly the simplest minimal affine Toda models which have particle species with different masses. We will compare numerical values of c_G^{eff} and c_g^{eff} again setting $g_{ab} = \frac{1}{G} \delta_{ab}$. For this purpose we use (49) and (69), taking into account [4] that the N -matrix in these equations is given by $N_{\mathbf{g}} = I_{\mathbf{g}}(2 - I_{\mathbf{g}})^{-1}$, where $I_{\mathbf{g}}$ stands for the incidence matrix of the corresponding Lie algebra $\mathbf{g}$.

In the A_3 case the masses of the three species are $m_1 = m_3 = m_2/\sqrt{2}$. Then, computing c_G^{eff} by formulae (49) and (52), we find: $c^{\text{eff}} \approx 0.77$, $c^{\text{eff}} = 1$, $c^{\text{eff}} \approx 1.12$, $c^{\text{eff}} \approx 1.16$ for $G = \frac{1}{2}, 1, 2, \infty$, respectively. The values of c_g^{eff} corresponding to the Haldane-Wu statistics with $g_{ab} = \frac{1}{G} \delta_{ab}$ were found in [23]: $c^{\text{eff}} \approx 0.89$, $c^{\text{eff}} = 1$, $c^{\text{eff}} \approx 1.07$, $c^{\text{eff}} \approx 1.16$ for $g = 2, 1, \frac{1}{2}, 0$, respectively. Carrying out analogous computations for the A_4 case, where the masses of the four species are $m_1 = m_4$, $m_2 = m_3 = m_1(\sqrt{5}+1)/2$, we find (Gentile statistics): $c^{\text{eff}} \approx 0.92$, $c^{\text{eff}} = 8/7$, $c^{\text{eff}} \approx 1.25$, $c^{\text{eff}} \approx 1.28$ for $G = \frac{1}{2}, 1, 2, \infty$, respectively; and (Haldane-Wu statistics): $c^{\text{eff}} \approx 1.05$, $c^{\text{eff}} = 8/7$, $c^{\text{eff}} \approx 1.20$, $c^{\text{eff}} \approx 1.28$ for $g = 2, 1, \frac{1}{2}, 0$, respectively.

One can also consider a folded algebra case with two different masses, namely $A_4^{(2)}$, where $m_2 = m_1(\sqrt{5}+1)/2$. In general, one assigns to the $A_{2n}^{(2)}$ minimal affine Toda model a tad-pole graph [4, 27] such that the corresponding incidence matrix differs from that of A_n only by 1 in the lower-right entry. Then one can check that x_a in (49) and y_a in (69) coincide with those for the A_{2n} case (a well-known fact for the fermionic statistics [4, 13]). Consequently, for any statistics we have $c^{\text{eff}}(A_{2n}^{(2)}) = \frac{1}{2} c^{\text{eff}}(A_{2n})$. Therefore, the data for $A_4^{(2)}$ follow from the above results for A_4 .

Finally, in the D_4 case, where the masses of the four species are $m_1=m_3=m_4=m_2/\sqrt{3}$, we obtain for the Gentile statistics: $c^{\text{eff}} \approx 0.82$, $c^{\text{eff}} = 1$, $c^{\text{eff}} \approx 1.08$, $c^{\text{eff}} \approx 1.09$ for $G = \frac{1}{2}, 1, 2, \infty$, respectively; and for the Haldane-Wu statistics: $c^{\text{eff}} \approx 0.93$, $c^{\text{eff}} = 1$, $c^{\text{eff}} \approx 1.04$, $c^{\text{eff}} \approx 1.09$ for $g = 2, 1, \frac{1}{2}, 0$, respectively.

We see that all the above results are in agreement with (68) and (72). This, together with the more detailed results we obtained in the A_1 and A_2 cases, allows us to conjecture that these inequalities and the inequality (63) for the scaling functions hold actually for all simply-laced minimal affine Toda models.

We remark also that in all the computations of c^{eff} in this subsection we always have the rule that the value of c_a is smaller for the species which has heavier particle (this was known for the fermionic statistics [4]). This observation provides an additional support to the above discussed interpretation of $\frac{6}{\pi^2}c_a$ as a massless degree of freedom of the corresponding particle.

8 Conclusion

Summarizing, we considered possible types of a generalized extensive statistics and established properties of the corresponding entropy densities. In particular, we established numerically (and gave some supporting analytical arguments) that the entropy density for the Gentile statistics of order $G > 1$ majorizes that for the Haldane-Wu statistics if they correspond to the same maximal occupancy. Further, we derived the thermodynamic Bethe ansatz equation and the finite-size scaling function $c(r)$ for a relativistic multi-particle system obeying a generalized extensive exclusion statistics. We put particular emphasis on the ultraviolet limit of such a system. We derived an expression for an effective central charge in the general case and showed that for the Gentile statistics it acquires an elegant form involving dilogarithms. We discussed a physical interpretation of the ‘partial’ central charges c_a and argued that it possibly leads to restrictions on the choice of statistics. Finally, we observed (and partially proved) that the majorizing properties of the Gentile statistics with respect to the Haldane-Wu statistics extend also to finite-size scaling functions and central charges related to the minimal simply-laced affine Toda models.

In the presented analysis of thermodynamic properties of a relativistic multi-particle system the single-state partition function $f(t)$ played a key role. For a given model, if we know $f(t)$ (at least in the thermodynamic limit) and the quantities $\varphi_{ab}(\theta)$, then we can compute, at least in principle, the entire scaling function which provides complete thermodynamic description of a system. In principle, $f(t)$ should be determinable from the corresponding Hamiltonian. However, this might be a non-trivial task since the exclusion statistics of particles can differ from the exchange statistics of the fields present in the Hamiltonian. Therefore, it may be more practical for a given system to fix an exotic statistics a-priori and investigate whether it leads to realistic physical properties. In particular, one can obtain information about the ultraviolet limit and see if it corresponds to an appropriate conformal field theory. As we have seen above, the different types (8) of $f(t)$ lead to significantly different thermodynamic properties. The question, which of these statistics can emerge in physical models, remains open.

Another open problem is to prove rigorously the assertion (27) that the entropy density for the Gentile statistics majorizes that for the Haldane-Wu statistics. Such a proof will probably provide a deeper insight into general properties of exotic exclusion statistics. Also

it would be interesting to verify the majorizing inequalities for the scaling functions (63) and central charges (68) related to the affine Toda models and to understand whether such a majorization is a model independent feature of the two involved statistics.

Acknowledgements: I am grateful to A. Fring for proposing the problem and participation in the initial stage of the research. This work was supported in part by INTAS grant 99-01459 and by Russian Fund for Fundamental Investigations grant 99-01-00101.

References

- [1] O.W. Greenberg and A.M.L. Messiah, *Phys. Rev.* **B136** (1964) 248.
- [2] F.D.M. Haldane, *Phys. Rev. Lett.* **67** (1991) 937.
- [3] F. Wilczek, *Phys. Rev. Lett.* **48** (1982) 1114, **49** (1982) 957;
Y.S. Wu, *Phys. Rev. Lett.* **52** (1984) 2103.
- [4] T.R. Klassen and E. Melzer, *Nucl. Phys.* **B338** (1990) 485.
- [5] K. Schoutens, *Phys. Rev. Lett.* **79** (1997) 2608;
A.G. Bytsko and A. Fring, *Nucl. Phys.* **B521** (1998) 573, *Phys. Lett.* **B454** (1999) 59;
P. Bouwknegt and K. Schoutens, *Nucl. Phys.* **B547** (1999) 501.
- [6] G. Gentile, *Nuovo Cim.* **17** (1940) 493; *Nuovo Cim.* **19** (1942) 109.
- [7] H.S. Green, *Phys. Rev.* **90** (1953) 270.
- [8] H.S. Green, *Prog. Theor. Phys.* **47** (1972) 1400;
Y. Ohnuki, S. Kamefuchi, *Prog. Theor. Phys.* **52** (1974) 1369;
R.Y. Levine, Y. Tomozawa, *Phys. Lett.* **B128** (1983) 189.
- [9] A.J. Bracken and H.S. Green, *J. Math. Phys.* **14** (1973) 1784;
P.G.O. Freund, *Phys. Rev.* **D13** (1976) 2322;
M. Cattani and N.C. Fernandes, *Nuovo Cim.* **B87** (1985) 70; *Phys. Lett.* **A124** (1987) 229.
- [10] J.D. Anand, S.N. Biswas and M. Hasan, *Phys. Rev.* **D18** (1978) 4529;
M.C. de Sousa Vieira and C. Tsallis, preprint CBPF-NF-038/86 (1986);
P. Suranyi, *Phys. Rev. Lett.* **65** (1990) 2329;
A.K. Rajagopal, *Phys. Lett.* **A214** (1996) 127.
- [11] K. Byczuk, J. Spalek, G.S. Joyce and S. Sarkar, *Acta Phys. Polon.* **B26** (1995) 2167.
- [12] C.N. Yang and C.P. Yang, *Phys. Rev.* **147** (1966) 303; *J. Math. Phys.* **10** (1969) 1115.
- [13] Al.B. Zamolodchikov, *Nucl. Phys.* **B342** (1990) 695; **B358** (1991) 524.
- [14] Al.B. Zamolodchikov, *Phys. Lett.* **B253** (1991) 391.
- [15] C. Nayak, F. Wilczek, *Phys. Rev. Lett.* **73** (1994) 2740.
- [16] K.N. Ilinski, J.M.F. Gunn and A.V. Ilinskaia, *Phys. Rev.* **B53** (1996) 2615.
- [17] A.P. Polychronakos, *Phys. Lett.* **B365** (1996) 202.
- [18] M.V. Fedoruk, *Asymptotics: integrals and series* (in Russian), (Nauka, 1987).
- [19] E.C. Titchmarsh, *The theory of functions*, (Oxford University Press, Oxford, 1939).
- [20] R. Acharya and P. Narayana Swamy, *J. Phys.* **A27** (1994) 7247;
A.V. Ilinskaia, K.N. Ilinski and J.M.F. Gunn, *Nucl. Phys.* **B458** (1996) 562.
- [21] Y.S. Wu, *Phys. Rev. Lett.* **73** (1994) 922.
- [22] D. Bernard and Y.S. Wu, cond-mat/9404025 (1994);
M.V.N. Murthy and R. Shankar, *Phys. Rev. Lett.* **73** (1994) 3331;
K. Hikami, *Phys. Lett.* **A205** (1995) 364;
S.B. Isakov, D.P. Arovas, J. Myrheim and A.P. Polychronakos, *Phys. Lett.* **A212** (1996) 299.
- [23] A.G. Bytsko and A. Fring, *Nucl. Phys.* **B532** (1998) 588.
- [24] A. Fring, C. Korff and B.J. Schulz, *Nucl. Phys.* **B549** (1999) 579.
- [25] C. Korff, hep-th/0008200 (2000).
- [26] H.W.J. Blöte, J.L. Cardy and M.P. Nightingale, *Phys. Rev. Lett.* **56** (1986) 742;
I. Affleck, *Phys. Rev. Lett.* **56** (1986) 746.
- [27] F. Ravanini, A. Valleriani and R. Tateo, *Int. J. Mod. Phys.* **A8** (1993) 1707;
M.J. Martins, *Nucl. Phys.* **B394** (1993) 339.
- [28] M.D. Freeman, *Phys. Lett.* **B261** (1991) 57;
A. Fring, H.C. Liao and D.I. Olive, *Phys. Lett.* **B266** (1991) 82.




Fig. 1: Entropy densities for $G = 2$ Gentile statistics (the upper curve), for $g = 1/2$ Haldane-Wu statistics (the middle curve) and for the fermionic statistics (the lower curve).




Fig. 2: Scaling functions of the A_1 -minimal affine Toda model for the Gentile statistics of order $G = 1, 2, 3, 4$ (solid curves, from down to up), for $g = 1/2$ Haldane-Wu statistics (the dashed curve) and for the bosonic statistics (the dotted curve).




Fig. 3: Solution $L(\theta)$ of the TBA equation for the A_2 -minimal affine Toda model for the scaling parameter $mr = 0.1$ (dotted curves), $mr = 0.01$ (dashed curves), and $mr = 0.001$ (solid curves). For the same value of mr the lower curve corresponds to the fermionic statistics and the upper curve to $G=2$ Gentile statistics.




Fig. 4: Scaling functions of the A_2 -minimal affine Toda model for the Gentile statistics of order $G = 1, 2, 3, 4$ (solid curves, from down to up) and for the Haldane-Wu statistics with $g_{ab} = \frac{1}{2}\delta_{ab}$ (the dashed curve).




Fig. 5: Effective central charge of the A_1 and A_2 -minimal affine Toda models for the Gentile statistics of order G (lower and upper solid curves) and for the Haldane-Wu statistics with $g_{ab} = \frac{1}{G}\delta_{ab}$ (lower and upper dashed curves).